RFERENCES:

1) Bostwick, D. G. (2002). "The Pathology of Benign Prostatic Hyperplasia". In Kirby, Peter; McConnell, John D.; Fitzpatrick, John M. et al.. Textbook of Benign Prostatic Hyperplasia. London: Isis Medical Media

2) Fisher R, Van Emde Boas W, Blume W, Elger C, Genton P, Lee P, Engel J (2005). "Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE)". Epilepsia 46 (4): 470–472.

3) "Epilepsy". World Health Organization. January 2009. http://www.who.int/mediacentre/factsheets/fs999/en/. Retrieved January 5, 2012

4) DIRC Newsletter Apr-Jun (2012) 14(2).,3.

5) Maryadele J. O' Neil (2001), THE MERCK INDEX, Merck Research Laboratories, Division of Merck & Co., Whitehouse Station, NJ, USA, 13th edition.,1679.

6) Sweetman S. C. Paul S B, Julie M .M. G (2005), The Complete Drug Reference, Pharamceutical Press., The Royal Pharmaceutical Society of Great Britain, 34th edition., 378.

7) Shrivastava A, Gupta V.B (2012), Stability-Indicating RP-HPLC Method for the Simultaneous Determination of Prazosin, Terazosin and Doxazosin in Pharmaceutical Formulations, Sci. Pharm, 80(3)., 619–631.

8) Kategaonkar A H, Patel D.M, Itkar B.P, Choudhari V.P, Kuchekar B. S and Nikalje A.G (2009), Spectrophotometric and Stability Indicating RP-HPLC-PDA method for Simultaneous Determination of Finasteride and Tamsulosin in Combined Tablet Dosage form, Asian Journal of research in chemistry,2(4)., 414-419.

9) Dhanya B, Suganthi A, Sen A, Sahoo U, Seth A. K (2011), Determination of Doxazosin Mesylate in Tablets by RP-HPLC, Indian J Pharm Sci, 73(1).,120–122.

10) Pavan B.A, Rama Krishna U.V, Vijayalakshmi M, Ravi Kumar A(2012), Spectrophotometric estimation of naftopidil in bulk and dosage form Res J Pharm. App Sci, 2(4).,26-30.

11) Pavan B. A, Vijayalakshmi M. And Manoj P (2012),Development And Validation Of Rp-Hplc Method For The Estimation Of Naftopidil In Bulk And Dosage Form, Ijrpc, 2(3).,816-21.

12) Dabhi B, Pandya D, Parmar A, Jadeja Y, Patel M , Jebaliya H, Karia D, Shah A (2012), Method Development And Validation Of Stability Indicating Isocratic Rp-Uplc Method For Assay Of Terazosin Hydrochloride Dihydrate In Pharmaceutical Tablets, Inventi Rapid: Pharm Analysis & Quality Assurance, Available From http://www.inventi.in/Article/ppaqa/453/12.aspx

13) Patel B, Patel J, Singh H, Patel B (2011), Extractive Spectrophotometric Methods for the Determination of gabapentin in Pharmaceutical Dosage Forms, IJPSDR, 3(3)., 197-201.

14) Cetin S. M, Atmaca S (2004),Determination of vigabatrin in human plasma and urine by high-performance liquid chromatography with UV-Vis detection, J.Chromato, 1031(1-2).,237-242.

15) Chang S.Y, Lin W C (2003),Determination of vigabatrin by capillary electrophoresis with laser-induced fluorescence detection, J.Chromato.B, 794(1), 17-22.

16) Sharma C R, Akhtar J, Jagani N M, Shankharva Y R, Shah J S (2012), Inventi Rapid: Pharm Analysis & Quality Assurance , Vol. 2012 , Article ID- " Inventi:ppaqa/310/12 " , [cited 2012 Sep 02] Available From http://www.inventi.in/Article/ppaqa/310/12.aspx

17) Bhavesh D, Pinal S, Saroj V, Shivprakash (2002), Determination of Doxazosin in human plasma by reversed phase ion-pair HPLC with fluorescence Detection, IJPS, 64(4)., 354-356.

18) Dhana S.M, Shaik R, Aravind. G, RP-HPLC Method For Quantitative Estimation Of Clobazam In Pharmaceutical Dosage Forms, IJRRPAS, 2(4).,751-759.

19) Patel D.B and Patel N.J, (2010), Validated RP-HPLC and TLC methods for simultaneous estimation of tamsulosin hydrochloride and finasteride in combined dosage forms, Acta Pharm. 60.,197–205.

20) Meghana D, Lahari K, Shantha Kumari K, Prakash K (2012), Development And Validation Of RP-HPLC Method For Simultaneous Estimation Of Clonazepam And Propranolol Hydrochloride In Bulk And Pharmaceutical Dosage Forms, Inventi Rapid: Pharm Analysis & Quality Assurance, 2012/9/17

21) Farnoush F, Ganjali M. R, Larijani B, Ensieh E Riahi S, Norouzi P (2010), Quantitative Analysis of Prazosin Hydrochloride in Pharmaceutical Formulation by Prazosin Potentiometric Sensor Based on Computational Investigation, Int. J. Electrochem. Sci, 5., 653 – 667.

22) Ramanaiah G, Ramachandran D, Srinivas G, Srilakshmi, Rao P (2012), Development and validation of UV spectroscopy method for estimation of lacosamide in bulk and formulations, Int J Biomed Sci,3(1), 10-12

23) Kumar G. S and B Sai Pavan Kumar (2012), Stability-Indicating RP-HPLC Method for Determination of Tamsulosin HCL in Pharmaceutical Dosage Form, Journal of Basic and Clinical Pharmacy, 3(2).,255-60.

24) Padma Latha H, Dr. Vidya S. G (2011), Estimation of Doxazocin Mesylate , an A-1- adrenergic receptor in Pharmaceutical dosage form by U.V.Spectrophotometric method, JPR,4(6)., 1642-43. ISSN: 0974-6943

25) Boda J. M, Bhalodiya H. A, Patel P. B (2012), Uv - spectroscopic method for simultaneous estimation of clonazepam and paroxetine hydrochloride hemihydrate in combined pharmaceutical formulation, Inventi Rapid: Pharm Analysis & Quality Assurance

26) Raghubabu K, Shanti swarup L, Kalyanaramu B, Rao M.N, Ramdas C (*2012),*Simple and Inexpensive Methods Development for the estimation of Tamsulosin Hydrochloride as a single component from Its Solid Dosage Forms by Visible spectrophotometry, IJPBS, 2(1).,12-19.

27) Linda E. Gustavson, Chu S (1992), High-Performance Liquid Chromatographic Procedure For The Determination Of Tiagabine Concentrations In Human Plasma Using Electrochemical Detection, J Chromatogr, 574(2).,313-8.
28) Thimmaraju M K, Rao V, Hemanth K, Siddartha Kumar P (2011), RP HPLC Method for the determination ofTamsulosin in bulk and Pharmaceutical Formulations, JAPS, 1(8).,177-180.
29) Naidu N.V.S, Reddy M. R. M and Suguna P (2012), Development And Validation Of The HPLC Method For The Analysis Of Doxazosin In Bulk Drug And Pharmaceutical Dosage Forms, IJPSR, 3(8)., 2705-2711.
30) Nevin O, Erturk S, Atmaca S (2002),Spectrofluorimetric and Spectrophotometric methods for the determination of vigabatrin in tablets, JPBA, 29(1-2)., 1-5.
31) Pande V.V, Jadhav J.N, Chandorkar J.G (2009), Estimation of Tamsulosin hydrochloride and Dutasteride in combined dosage form by UV spectroscopy method, Journal.P. Res, 2(3)., 507-509.
32) Gur P, Poklis A ,Joseph S (1995), Chromatographic Procedures for the Determination of Felbamate in Serum, J Anal Toxicol, 19 (6)., 499-503.
33) Satyanarayana PVV, and Alavala S M (2012), Validated RP - HPLC Method for the Estimation of Ezogabine in Tablet Dosage Form,I.J.R.P.B.S, 3 (2)., 955-59.
34) Jat R. K, Chhip R. C and Sharma S (2010), Quantitative Estimation of Clobazam in Bulk drug and tablets, I.J.C.P.R,1(3)., 18-24.
35) Gujral R. S, Manirul H, Prem S. K (2010), A Novel Quantitative Spectrometric Method for the Analysis of Gabapentin Hydrochloride, J.P.S.Tech,2(5)., 222-229.
36) Richa Kumari, Dash P. P, Lal V. K, Mishra A , and Murthy P. N (2010) ,RP – HPLC method for the estimation of Tamsulosin Hydrochloride in Tablet Dosage Form, Indian J Pharm Sci, 72(6).,785–787.
37) Rustum AM, Menon G, Patel S (1998), Separation of the S(+) and R(-)-enantiomers of tiagabine.HCl and its two chiral precursors by chiral chromatography: application to chiral inversion studies, J Pharm Biomed Anal, 17(8).,1439-47.

38) Kamdar S.A, Vaghela V. M, Desai P. A (2012), Development and Validation of HPTLC Method for Estimation of Lacosamide in Bulk Drug and in Tablet Dosage Form, I.J.C.R.G.G, 4(3)., 1193-1197.

39) Sawant D. S,Ghante M. R, Deshpande A. S, Munot N. M (2012),Devolopment and Validation of RP-HPLC Method for the Estimation of Tamsulosin Hydrochloride in Bulk Drug and Pharmaceutical Dosage Form, J P R, 3(11).,2776-2778.

40) Shabana Naz, Najma Sultana, Saeed M. A, Nighat Shafi (2010), Simultaneous Determination Of Prazosin And Calcium Chanel Blockers In RawMaterials, Pharmaceutical Formulations And Human Serum By Rp-Hplc,I.J.P.R.D, 2(9).,6-16.

41) Usmangani K. Chhalotiya Kashyap K. Bhatt D. A, Sha S. L, Baldania J, Patel R (2012), Stability-Indicating Liquid Chromatographic Method For Quantification Of New Anti-Epileptic Drug Lacosamide In Bulk And Pharmaceutical Formulation Cheml Indus & Cheml Eng Quarterly, 18 (1)., 35-42.

42) Chakravarthy V. K and Gowri Sankar D (2011), Development and Validation of RP-HPLC method for estimation of lacosamide in bulk and its pharmaceutica formulation, 4(3)., 666-672.

43) Galande V. R , Baheti K.G , Dehghan M. H (2010), UV-Vis Spectrophotometric Method For Estimation Of Gabapentin Methylcobalamin In Bulk And Tablet, I.J.C.R.G.G, 2(1)., 695-699.

44) Patel V. B , Dave J.B, Patel F.M, Chhaganbhai N (2012), Uv Spectrophotometric Method For Identification And Estimation Of Clonazepam In Tablet Dosage Form, 1(2)., 62-70.

45) Xia Z, Liu Y, Xu J, Zhang D, Zhou Y, Gu J, Cui Y (2009), Determination of Silodosin in human plasma by Lliquid Chromatography- Tandom Mass Spectroscopy, J Chromatogr B Analyt Technol Biomed Life Sci,877(29).,3724-8.

46) Sun Y. S , Huang B. Y, Mu Yuan , Shi J. S (2009), Development of a chiral HPLC method for the analysis of naftopidil enantiomers, JCPS 18 (1)., 61-63.